

OBSERVACIONES	ORNAMENTALES (SL)	Moho gris (<i>Botrytis cinerea</i>) Mancha negra (<i>Diplocarpum rosae</i>) Cenicilla (<i>Sphaerotheca pannosa</i>) Tizón tardío (<i>Phytophthora infestans</i>) Tizón temprano (<i>Alternaria solani</i>) Damping off, Pudrición radical (<i>Rhizoctonia sp.</i>) Pudrición radical (<i>Fusarium sp.</i>) Mancha café de la hoja (<i>Cercospora arachidicola</i>) Mildiu (<i>Pseudoperonospora cubensis</i>) Antracnosis (<i>Colletotrichum lagenarium</i>) Podredumbre gomosa de los tallos (<i>Mycosphaerella citrulina</i>) Tizón foliar (<i>Alternaria cucumerina</i>) Mancha foliar (<i>Cercospora cassicola</i>) Mancha foliar (<i>Cercospora sp.</i>) Cenicilla (<i>Erysiphe cichoracearum</i>) Roña (<i>Cladosporium cucumerinum</i>) Mancha negra (<i>Diplocarpum rosae</i>) Moho gris (<i>Botrytis cinerea</i>) Mildiu (<i>Pseudoperonospora cubensis</i>) Tizón foliar (<i>Alternaria cucumerina</i>) Mancha foliar (<i>Corynespora cassicola</i>) Podredumbre gomosa de los tallos (<i>Cercospora sp.</i>) Podredumbre gomosa de los tallos (<i>Mycosphaerella citrulina</i>) Antracnosis (<i>Colletotrichum lagenarium</i>) Roña (<i>Cladosporium cucumerinum</i>) Mancha ojo de rana (<i>Cercospora sojae</i>) Mancha foliar (<i>Cercospora sp.</i>) Antracnosis (<i>Colletotrichum dematium</i>) Antracnosis (<i>Colletotrichum lindemuthianum</i>) Mildiu (<i>Pseudoperonospora cubensis</i>) Tizón de la vaina y tallo (<i>Diaporthe phaseolorum</i>) Mancha foliar (<i>Sclerotinia sp.</i>) Tizón temprano (<i>Alternaria solani</i>) Mancha púrpura (<i>Cercospora kikuchii</i>) Antracnosis (<i>Colletotrichum sp.</i>) Antracnosis (<i>Colletotrichum phomoides</i>) Tizón temprano (<i>Alternaria solani</i>) Tizón tardío (<i>Phytophthora infestans</i>) Tizón foliar (<i>Cercospora carotae</i>) Tizón foliar (<i>Alternaria dauci</i>)	1.25-1.7 L / 100 L de agua 1.2-2.5 L/ha 0.9-2.5 L/ha 1.5-2.0 L/ha 1.2-1.7 L/ha 1.5-3.50 L/ha 1.5-2.50 L/ha 1.5-3.5 L/ha 1.5-3.5 L/ha 1.5-3.5 L/ha 1.5-3.5 L/ha 2.5 L/ha 2.5-3.5 L/ha 135-170ml/ 100 L de agua 1.50-3.50 L/ha 1.50-2.50 L/ha 1.50-3.50 L/ha 1.50-2.50 L/ha 2.5-3.5 L/ha 2.5-3.5 L/ha 1.5-2.5 L/ha 1.2-2.5 L/ha 1.5-2.5 L/ha 1.2-2.5 L/ha 1.20-2.50 L/ha 1.2-2.5 L/ha 1.2-3.0 L/ha 1.5-2.5 L/ha 2.5-3.5 L/ha 1.75-3.5 L/ha 1.50-2.50 L/ha 2.5 L/ha 3.5-5.0 L/ha 350-450 mL/100 L de agua 3.5-5.0 L/ha 150-450 mL/ 100 L de agua	OBSERVACIONES	Realizar la primera aplicación de manera preventiva cuando las condiciones climáticas sean favorables para el desarrollo de la plaga.
1.75-2.5 L/ha 1.75 L/ha 1.75 L/ha 1.75-2.5 L/ha 1.75 L/ha	PAPA (SL)				
135-170 mL/ 100 L agua 3.5-5.0 L/ha	PEPINO (SL)		Aplicación foliar		
350-450 mL/100 L de agua 2.5-3.0 L/ha 2.5 L/ha	PEPINO (SL)				
3.5 L/ha 3.5-5.0 L/ha 350-450 mL/100 L de agua 3.5-5.0 L/ha	ROSAL (SL)				
35-170 mL/ 100 L de agua 1.5-2.5 L/ha	SANDÍA (SL)				
1.5-3.5 L/ha 1.5-3.5 L/ha	SOYA (42)			Aplicación foliar	
35-170 mL/ 100 L de agua 0.875-2.5 L/ha	TOMATE DE CASCARA (SL)				
1.50-3.50 L/ha 2.5-3.5 L/ha 1.75-3.5 L/ha 1.50-2.50 L/ha 2.5 L/ha 3.5-5.0 L/ha	ZANAHORIA (SL)				
350-450 mL/100 L de agua 3.5-5.0 L/ha 150-450 mL/ 100 L de agua	OBSERVACIONES				

CULTIVO	PLAGA	DOSIS	OBSERVACIONES	COL DE BRUSELAS (SL)	Mancha foliar (<i>Alternaria brassicae</i>) Mildiu (<i>Peronospora parasitica</i>)
AJO (14)	Mancha púrpura (<i>Alternaria porri</i>)	1.5-3.5 L/ha		COLIFLOR (SL)	Mancha anular (<i>Mycosphaerella brassicicola</i>)
	Moho gris (<i>Botrytis allii</i>)	1.2-3.5 L/ha			Mildiu (<i>Peronospora parasitica</i>)
	Moho gris (<i>Cladosporium fulvum</i>)	1.20-2.5 L/ha			Mancha anular (<i>Mycosphaerella brassicicola</i>)
	Mildiu (<i>Peronospora destructor</i>)	1.2-3.5 L/ha			Moho gris (<i>Botrytis</i> sp.)
	Mancha foliar (<i>Corynespora cassicola</i>)	1.20-2.5 L/ha			Mancha foliar (<i>Cercospora chrysanthemi</i>)
APIO (7)	Mancha foliar (<i>Cercospora apii</i>)	1.2-1.75 L/ha		CRISANTEMO (SL)	Mancha foliar (<i>Cercospora chrysanthemi</i>)
	Putridión del tallo y raíz (<i>Sclerotinia</i> sp.)	1.20-1.75 L/ha			Verrucosis (<i>Taphrina deformans</i>)
	Mancha foliar o tizón (<i>Septoria apii</i>)	1.2-3.5 L/ha			Putridión morena (<i>Monilinia fructicola</i>)
	Damping off, Putridión radical (<i>Rhizoctonia</i> sp.)	1.5-3.5 L/ha			Cenicilla (<i>Sphaerotheca humuli</i>)
	Roya (<i>Puccinia arachidis</i>)				Tizón tardío (<i>Phytophthora infestans</i>)
CACAHUATE (14)	Mancha café de la hoja (<i>Cercospora arachidicola</i>)	1.2-1.75 L/ha		CHABACANO (30)	Tizón temprano (<i>Alternaria solani</i>)
	Tizón temprano (<i>Alternaria solani</i>)				Mancha foliar (<i>Cercospora capsici</i>)
	Mancha o peca de la hoja (<i>Cercospora personata</i>)				Tizón foliar (<i>Alternaria dauci</i>)
	Mal de hilachas (<i>Corticium kolerogae</i>)				Tizón foliar (<i>Alternaria alternata</i>)
	Putridión negra (<i>Phytophthora palmivora</i>)				Verrucosis (<i>Taphrina deformans</i>)
CACAO (SL)	Putridión del fruto (<i>Monilinia</i> sp.)	3-6 mL/planta		DURAZNO (14)	Tiro de munición (<i>Coryneum beijerinckii</i>)
	Tizón foliar (<i>Alternaria cucumerina</i>)	1.50-2.50 L/ha			Cenicilla (<i>Sphaerotheca humuli</i>)
	Roña (<i>Cladosporium cucumerinum</i>)	2.5-3.5 L/ha			Tizón tardío (<i>Phytophthora infestans</i>)
	Antracnosis (<i>Colletotrichum lagenarium</i>)				Tizón temprano (<i>Alternaria solani</i>)
	Mancha foliar (<i>Cercospora</i> sp.)	1.50-2.50 L/ha			Mancha foliar (<i>Cercospora capsici</i>)
CALABACITA (SL)	Mancha foliar (<i>Cercospora</i> sp.)	2.5 L/ha		CULTIVO	Tizón foliar (<i>Alternaria dauci</i>)
	Cenicilla (<i>Sphaerotheca humuli</i>)	1.75-3.5 L/ha			Tizón foliar (<i>Alternaria alternata</i>)
	Cenicilla (<i>Erysiphe cichoracearum</i>)	2.5-3.5 L/ha			Verrucosis (<i>Taphrina deformans</i>)
	Mildiu (<i>Pseudoperonospora cubensis</i>)				Tiro de munición (<i>Coryneum beijerinckii</i>)
	Podredumbre gomosa de los tallos (<i>Mycosphaerella citrulina</i>)				Putridión morena (<i>Monilinia fructicola</i>)
CALABAZA (SL)	Mildiu (<i>Pseudoperonospora cubensis</i>)	1.50-2.50 L/ha		DURAZNO (14)	Putridión morena (<i>Monilinia fructicola</i>)
	Antracnosis (<i>Colletotrichum lagenarium</i>)				Moho gris (<i>Botrytis cinerea</i>)
	Tizón foliar (<i>Alternaria cucumerina</i>)	2.5 L/ha			Antracnosis (<i>Colletotrichum lindemuthianum</i>)
	Mancha foliar (<i>Corynespora cassicola</i>)	1.75-3.5 L/ha			Roya (<i>Uromyces appendiculatus</i>)
	Mancha foliar (<i>Cercospora</i> sp.)	2.5-3.5 L/ha			Mildiu (<i>Peronospora parasitica</i>)
CEBOLLA VERDE (14) CEMPASUCHIL (SL)	Mancha foliar (<i>Cercospora</i> sp.)	1.50-2.50 L/ha		FRUJOL (SL)	Putridión morena (<i>Monilinia fructicola</i>)
	Roña (<i>Cladosporium cucumerinum</i>)				Moho gris (<i>Botrytis cinerea</i>)
	Cenicilla (<i>Erysiphe cichoracearum</i>)				Roya (<i>Uromyces appendiculatus</i>)
	Mancha púrpura (<i>Alternaria porri</i>)	1.75-3.5 L/ha			Moho gris (<i>Botrytis</i> sp.)
	Mancha foliar (<i>Botrytis</i> sp.)	1.5-2.5 L/ha			Roya (<i>Uromyces phaseoli</i>)
CÉSPED (SL)	Mancha foliar (<i>Alternaria</i> sp.)	125-250 mL/100 m ²	Aplicación foliar	MAÍZ (14)	Moho gris (<i>Botrytis cinerea</i>)
	Mancha foliar (<i>Alternaria</i> sp.)	200-250 mL/100 m ²			Roya (<i>Puccinia graminis</i>)
	Mancha foliar (<i>Alternaria</i> sp.)	60-120 mL/100 m ²			Tizón foliar (<i>Helminthosporium maydis</i>)
	Putridión radical (<i>Rhizoctonia solani</i>)	125-200 mL/100 m ²			Tizón foliar o mancha foliar (<i>Helminthosporium sp.</i>)
	Moho gris (<i>Botrytis</i> sp.)	3.5-5.0 L/ha			Mancha foliar (<i>Corynespora cassicola</i>)
CIRUELO (14)	Mancha foliar (<i>Puccinia griseae</i>)			MELÓN (SL)	Podredumbre gomosa de los tallos (<i>Mycosphaerella ol</i>)
	Putridión radical (<i>Rhizoctonia solani</i>)	350-450 mL/100L de agua			Tizón foliar (<i>Alternaria cucumerina</i>)
	Moho gris (<i>Botrytis</i> sp.)	3.5-5.0 L/ha			Roña (<i>Cladosporium cucumerinum</i>)
	Tizón o mancha foliar (<i>Helminthosporium</i> sp.)				Mildiu (<i>Pseudoperonospora cubensis</i>)
	Roya (<i>Puccinia graminis</i>)				Antracnosis (<i>Colletotrichum lagenarium</i>)
COL (SL)	Verrucosis (<i>Taphrina deformans</i>)	1.75-2.5 L/ha		NECTARINO (30)	Cenicilla (<i>Sphaerotheca humuli</i>)
	Mancha foliar (<i>Cercospora</i> sp.)				Verrucosis (<i>Taphrina deformans</i>)
	Mancha foliar (<i>Alternaria brassicae</i>)				Tiro de munición (<i>Coryneum beijerinckii</i>)
	Mildiu (<i>Peronospora parasitica</i>)				Gomosis, tiro de munición (<i>Cladosporium carpophyllum</i>)
	Mancha anular (<i>Mycosphaerella brassicicola</i>)				Putridión morena (<i>Monilinia fructicola</i>)